

DENSITY IS AT MOST THE SPREAD OF THE SQUARE

SAHARON SHELAH

The Hebrew University of Jerusalem
Einstein Institute of Mathematics
Edmond J. Safra Campus, Givat Ram
Jerusalem 91904, Israel

Department of Mathematics
Hill Center-Busch Campus
Rutgers, The State University of New Jersey
110 Frelinghuysen Road
Piscataway, NJ 08854-8019 USA

I would like to thank Alice Leonhardt for the beautiful typing.
Publication E56

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\text{\texttt{TeX}}$

§1

1.1 Claim. Assume \mathbb{B} is an infinite Boolean Algebra and $\lambda = d(\mathbb{B})$. Then $\mathfrak{s}(\mathbb{B} * \mathbb{B})$, i.e. $\mathfrak{s}(\text{uf}(\mathbb{B}) \times \text{uf}(\mathbb{B})) \geq \lambda$ (if λ limit-obtained).

Remark. 1) $\text{ul}(\mathbb{B})$ is the space of ultrafilters of \mathbb{B} a compact space with clopen base.
 2) $\mathfrak{s}(X)$ is $\sup\{|Y| : Y \subseteq X \text{ is discrete } Y, \text{ same as } \text{des}(X)\}$.
 3) We meant to consider whether this works for compact Hausdorff spaces. But subsequently and independently Szentmiklössy prove this.

Proof. Without loss of generality $\lambda > \aleph_0$. We choose (p_i^0, p_i^1, a_i) by induction on $i < \lambda$ such that

- ⊗ (a) p_i^ℓ is an ultrafilter of \mathbb{B} for $\ell = 0, 1$
- (b) $a_i \in p_i^1, a_i \notin p_i^0$, i.e. $(-a_i) \in p_i^1$
- (c) if $j < i$ then $a_j \in p_i^0 \Leftrightarrow a_j \in p_i^1$
- (d) if $j < i$ then $a_i \notin p_j^0, a_i \notin p_j^1$.

So assume we have arrived to i . Let \mathbb{B}_i be the subalgebra of \mathbb{B} generated by $\{a_j : j < i\}$.

For every non-zero $b \in \mathbb{B}_i$ choose an ultrafilter q_b^i of \mathbb{B} and for simplicity $b = a_j \Rightarrow q_b^i = p_j^1$ and $b = (-a_j) \Rightarrow q_b^i = p_j^0$ for $j < i$.

As $d(\mathbb{B}) \geq \lambda$ clearly $\{q_b^i : b \in \mathbb{B}_i \setminus \{0\}\}$ is not dense hence there is a non-zero $a_i \in \mathbb{B}$ such that $b \in \mathbb{B}_i \setminus \{0\} \Rightarrow a_i \notin q_b^i$ (i.e. a non-empty clopen set to which none of the points q_b^i belongs).

Now clearly $b \in \mathbb{B}_i \setminus \{0\} \Rightarrow a_i \neq b$ (as $b \in q_p^i$) hence $a_i \notin \mathbb{B}_i$. This implies that there is an ultrafilter q_i^* of \mathbb{B}_i such that

$$\otimes \quad b \in q \Rightarrow a_i \cap b > 0 \wedge (-a_i) \cap b > 0.$$

[Why? As $\{b_0 \cup b_1 : b_0, b_1 \in \mathbb{B}_2 \text{ and } b - 1 \cap a_i = 0_{\mathbb{B}} \text{ and } b_a - a_i = 0\}$ is a proper ideal of \mathbb{B}_i hence can be extended to an ultrafilter of \mathbb{B}_i .]

So there are ultrafilters p_i^0, p_i^1 of \mathbb{B} such that

$$\otimes \quad q_i^* \cup \{a_i\} \subseteq p_i^1 \text{ and } q_i^* \cup \{-a_i\} \subseteq p_i^0.$$

This is enough for the induction step.

Having carried the induction

- (a) $p_i := (p_i^0, p_i^1) \in \text{uf}(\mathbb{B}) \times \text{uf}(\mathbb{B})$
- (b) $(-a_i) \times a_i$ is an open subset of $\text{uf}(\mathbb{B}) \times \text{uf}(\mathbb{B})$.

Lastly,

- (c) if $i < j < \lambda$ then $p_j \notin (-a_i) \times a_i$ because $(-a_i) \notin p_j^0$ or $a_i \notin p_j^1$ as $a_i \in \mathbb{B}_j$ and $p_j^0 \cap B_j = p_j^1 \cap \mathbb{B}_j$ by the choice of p_j^0, p_0^1
- (d) if $i < j < \lambda$ then $p_i \notin (-a_j) \times a_j$ because $p_i^1 \notin a_j$ by the choice of a_j .

So $\langle (p_i, (-a_i) \times a_i) : i < \lambda \rangle$ exemplify $\text{dis}(\text{uf}(\mathbb{B}) \times \text{uf}(\mathbb{B})) \geq \lambda$ as required.